

- [6] Wild-Altamirano C. Enzymic activity during growth of vanilla fruit. 1. Proteinase, glucosidase, peroxidase and polyphenoloxidase[J]. Journal of Food Science, 1969, 34: 235-238.
- [7] Arana F. E. Action of α -glucosidase in the curing of vanilla [J]. Food Research, 1943, 288: 343-351.
- [8] Odoux E., Chauwin A., Brillouet J. M. Purification and characterization of vanilla bean (*Vanilla planifolia* Andrews)-D-glucosidase[J]. Journal of Agricultural and Food Chemistry, 2003a, 51: 3168-3173.
- [9] Dignum M. J. W., Van Der Heijden R., Kerler J., Winkel C., Verpoorte R. Identification of glucosides in green beans of *Vanilla planifolia* Andrews and kinetics of vanilla-glucosidase[J]. Food Chemistry, 2004, 85: 199-205.
- [10] Odoux E., Escoute J., Verdeil J.-L., Brillouet J.-M. Localization of beta-D-glucosidase activity and glucovanillin in vanilla bean (*Vanilla planifolia* Andrews)[J]. Annals of Botany, 2003, 92: 437-444.
- [11] Odoux E., Escoute J., Verdeil J.-L. The relation between glucovanillin, beta-D-glucosidase activity and cellular compartmentation during the senescence, freezing and traditional curing of vanilla beans[J]. Annals of Applied Biology, 2006, 149: 43-52.
- [12] 江明, 浦帆. 香荚兰成熟荚果中三种酶的活性测定[J]. 云南植物研究, 2000a, 22(2): 187-190.
- [13] 江明, 刘涛. 不同加工处理条件下香荚兰荚果中二种内源酶的活性变化[J]. 云南植物研究, 2005, 27(3): 310-314.
- [14] Ranadive A S. Vanilla and related flavor compound in Vanilla extracts made from beans of various global origins[J]. J. Agric. Food Chem., 1992, 40: 1922-1924.
- [15] 浦帆, 江明. 香荚兰酶促生香的研究[J]. 云南植物研究, 1998a, 20(3): 355-361.
- [16] 浦帆, 张劲松. 不同阶段香荚兰豆荚中糖甙水解成份的研究[J]. 天然产物研究与开发, 1998b, 10(3): 29-33.
- [17] Odoux E. Changes in vanillin and glucovanillin concentrations during the various stages of the process traditionally used for curing *Vanilla fragrans* beans in Reunion[J]. Editions scientifiques et medicales Elsevier SAS, 2000, 55: 119-125.