

参考文献

- [7] Hernan J. C., Winniford B., Luong J., Matthias P. Comprehensive two dimensional gas chromatography review[J]. *J. Separation Sci.*, 2009, 32(5): 883-904.
- [8] Dallüge J., Beens J. Comprehensive two-dimensional gas chromatography: a powerful and versatile analytical tool[J]. *J. Chromatogr. A*, 2003, 1000(1): 69-108.
- [9] Rachel E. M., Kenneth M. D., Jamin C. H., Elton T. Y. Comprehensive two-dimensional gas chromatography time-of-flight mass spectrometry analysis of metabolites in fermenting and respiring yeast cells[J]. *Anal. Chem.*, 2006, 78(8): 2 700-2 709.
- [10] Karisa M. P., Janiece L. H., Jamin C. H., Synovec R. E. A principal component analysis based method to discover chemical differences in comprehensive two-dimensional gas chromatography with time-of-flight mass spectrometry (GC × GC-TOF-MS) separations of metabolites in plant samples[J]. *Talanta*, 2006, 70(4): 797-804.
- [11] Sanchez J. M., Sacks R. D. Development of a multibed adsorption trap, comprehensive two-dimensional gas chromatography, and time-of-flight mass spectrometry system for the analysis of volatile organic compounds in human breath[J]. *Anal. Chem.*, 2006, 78(9): 3 046-3 054.
- [12] Ognjen P., Tadeusz G., Comprehensive two-dimensional gas chromatography (GC × GC) in environmental analysis and monitoring[J]. *Anal. Bioanal. Chem.*, 2006, 386(4): 1 013-1 023.
- [13] Bester K. Analysis of musk fragrances in environmental samples[J]. *J. Chromatogr. A*, 2009, 1216(3): 470-480.
- [14] 汪秋安. 中国古近代香料史初探[J]. *香料香精化妆品*, 1999(2): 36-39.
- [15] Clark T. J., Bunch J. E. Qualitative and quantitative analysis of flavor additives on tobacco products using SPME-GC/MS[J]. *J. Agric. Food Chem.*, 1997, 45: 844-849.
- [16] O'zel M. Z., Gogus F., Lewis A.C., Subcritical water extraction of essential oils from *thymbra spicata*[J]. *Food Chem.*, 2003, 82(3): 381-386.
- [17] Zini C.A., Deassis T.F., Ledford E.B., Dariva C., Fachel J. Correlations between pulp properties of eucalyptus clones and leaf volatiles using automated solid-phase microextraction[J]. *J. Agric. Food Chem.*, 2003, 51: 7 848-7 853.
- [18] Davis, J. M., Giddings, J. C. Application to environmental chemistry. PAHs represent a class of organic pollutants that are widely found in the environment[J]. *Anal. Chem.*, 1983, 55: 418-424.
- [19] Giddings J. C., H. J. Multidimensional chromatography[M]. New York: Marcel Dekker, 1990, 1-27.
- [20] Cortes H. J. Developments in multidimensional separation systems[J]. *J. Chromatogr.*, 1992, 626: 3-23.
- [21] Liu Z., Sirimanne S. R., Patterson D. G., et al. Comprehensive two-dimensional gas chromatography for the fast separation and determination of pesticides extracted from human serum[J]. *Anal. Chem.*, 1994, 66: 3 086-3 092.
- [22] Phillips J. B., Ledford E. B. Thermal modulation: a chemical instrumentation component of potential value in improving portability[J]. *Field Anal. Chem. Tech.*, 1996, 1: 23-29.
- [23] Marriott P. J., Kinghom R. M. Longitudinally Modulated cryogenic system. a generally applicable approach to solute trapping and mobilization in gas chromatography[J]. *Anal. Chem.*, 1997, 69: 2 582-2 588.
- [24] Seeley J. V., Kramp F., Hicks C. J. Comprehensive two-dimensional gas chromatography via differential flow modulation[J]. *Anal. Chem.*, 2000, 72: 4 346-4 352.
- [25] Adahchour M., Beens J., Vreuls R. R., et al. Recent developments in comprehensive two-dimensional gas chromatography (GC × GC): I. introduction and instrumental set-up[J]. *Trends Anal. Chem.*, 2006, 25: 438-454.
- [26] Seeley J. V., Primeau N. J., Bandurski S. V., et al. Microfluidic Deans switch for comprehensive two-dimensional gas chromatography[J]. *Anal. Chem.*, 2007, 79: 1 840-1 847.
- [27] Poliak M., Kochman M., Amirav A. Pulsed flow modulation comprehensive two dimensional gas chromatography[J]. *J. Chromatogr. A*, 2008, 1 186: 189-195.
- [28] Frior R. Application Brief: Comprehensive GC system based on flow modulation for the 7890A GC[M]. Santa Clara: Agilent Technologies., 2008.
- [29] Adahchour M., Beens J., Vreuls R. J. J., Brinkman U.A. Comprehensive two-dimensional gas chromatography with time-of-flight mass spectrometric detection for the trace analysis of flavour compounds in food[J]. *J. Chromatogr. A*, 2003, 1019: 157-172.
- [30] Adahchour M., Beens J., Vreuls R. J. J., Batenburg A. M. Comprehensive two-dimensional gas chromatography of complex samples by using a'reversed-type' column combination: application to food analysis[J]. *J. Chromatogr. A*, 2004, 1 054: 47-55.
- [31] Zhu S., Lu X., Ji K., et al. Characterization of flavor compounds in Chinese liquor Moutai by comprehensive two-dimensional gas chromatography/time-of-flight mass spectrometry[J]. *Anal. Chim. Acta.*, 2007, 597: 340-348.
- [32] Eyres G. T., Marriott P. J., Dufour J. P. Comparison of odor-active compounds in the spicy fraction of hop (*Humulus lupulus* L.) essential oil from four different varieties[J]. *J. Agric. Food Chem.*, 2007, 55: 6 252-6 261.
- [33] Rochat S., Desaint L. J. Y., Chaintreau A. Analysis of sulfur compounds from the in-oven roast beef aroma by comprehensive two-dimensional gas chromatography[J]. *J. Chromatogr. A*, 2007, 1 147: 85-94.
- [34] d'Acampora Z. B., Casilli A., Dugo P., et al. Odour fingerprint acquisition by means of comprehensive two-dimensional gas chromatography-olfactometry and comprehensive two-dimensional gas chromatography/mass spectrometry[J]. *J. Chromatogr. A*, 2007, 1 141: 279-286.
- [35] Cardeal Z. L., Desouza P. P., Marriott P. J. Comprehensive two-dimensional gas chromatography for fingerprint pattern recognition in cachaca production[J]. *Talanta*, 2008, 74: 793-799.
- [36] Bianchi F., Careri M., Conti C., et al. Comparison of comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry and gas chromatography-mass spectrometry for the qualitative characterisation of roasted barley by solid-phase microextraction[J]. *J. Sep. Sci.*, 2007, 30: 527-533.
- [37] Cajka T., Hajslova J., Cochran J., et al. Solid phase microextraction-comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry for the analysis of honey volatiles[J]. *J. Sep. Sci.*, 2007, 30: 534-546.
- [38] Tran T. C., Marriott P. J. Characterization of incense smoke by solid phase microextraction-Comprehensive two-dimensional gas chromatography (GC × GC)[J]. *Atmos. Environ.*, 2007, 41: 5 756-5 768.

- [39] Vonmuhlen C., Zini C. A., Caramao E. B., et al. Comparative study of eucalyptus dunnii volatile oil composition using comprehensive two-dimensional gas chromatography coupled to time-of-flight and quadrupole mass spectrometry[J]. J. Chromatogr. A, 2008, 1 200: 34-42.
- [40] Eyres G., Marriott P. J., Dufour J. P. The combination of gas chromatography-olfactometry and multidimensional gas chromatography for the characterisation of essential oils[J]. J. Chromatogr. A, 2007, 1 150: 70-77.
- [41] Cordero C., Bicchi C., Galli M., et al. Evaluation of different internal-diameter column combinations in comprehensive two-dimensional gas chromatography in flavour and fragrance analysis[J]. J. Sep. Sci., 2008, 31: 3 437-3 450.
- [42] Shellie R.A., Marriott P. J., Chaintreau A. Quantitation of suspected allergens in fragrances (Part I): Evaluation of comprehensive twodimensional gas chromatography for quality control[J]. J. Flavour Frag., 2004, 19: 91-98.
- [43] Adahchour M., Brandt M., Baier H. U., et al. Comprehensive two-dimensional gas chromatography coupled to a rapid-scanning quadrupole mass spectrometer: principles and applications[J]. J. Chromatogr. A, 2005, 1 067: 245-254.
- [44] Zellner B. A., Casilli A., Dugo P., et al. Odour fingerprint acquisition by means of comprehensive two-dimensional gas chromatography-olfactometry and comprehensive two-dimensional gas chromatography/mass spectrometry[J]. J. Chromatogr. A, 2007, 1 141: 279-286.
- [45] Jover E., Adahchour M., Bayona J. M., et al. Characterization of lipids in complex samples using comprehensive two-dimensional gas chromatography with time-of-flight mass spectrometry[J]. J. Chromatogr. A, 2005, 1 086: 2-21.
- [46] O'zel M. Z., Gogus F., Hamilton J. F., et al. The essential oil of *pistacia vera* L. at various temperatures of direct thermal desorption using comprehensive gas chromatography coupled with time-of-flight mass spectrometry[J]. Chromatographia, 2004, 60: 79-83.
- [47] Roberts M. T., Dufour J. P., Lewis A. C. Application of comprehensive multidimensional gas chromatography combined with time-of-flight mass spectrometry (GC × GC-TOF/MS) for high resolution analysis of hop essential oil[J]. J. Sep. Sci., 2004, 27: 473-478.
- [48] Eyres G., Dufour J. P., Hallifax G., et al. Identification of character-impact odorants in coriander and wild coriander leaves using gas chromatography-olfactometry (GC-O) and comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry (GC × GC-TOF/MS)[J]. J. Sep. Sci., 2005, 28: 1 061-1 074.
- [49] Wu J. F., Lu X., Tang W. Y., et al. Application of comprehensive two-dimensional gas chromatography/time-of-flight mass spectrometry to analysis of zedoary volatile oil[J]. Chem. J. Chin. Univ., 2004, 25: 1 432-1 437.
- [50] Zhu S. H., Lu X., Dong L., et al. Quantitative determination of compounds in tobacco essential oils by comprehensive two-dimensional gas chromatography coupled to time-of-flight mass spectrometry[J]. J. Chromatogr. A, 2005, 1 086: 107-114.
- [51] Ma C. H., Wang H., Lu X., et al. Analysis of Artemisia annua L. volatile oil by comprehensive two-dimensional gas chromatography time-of-flight mass spectrometry[J]. J. Chromatogr. A, 2007, 1 150: 50-53.
- [52] Qiu Y., Lu X., Pang T., et al. Study of traditional Chinese medicine volatile oils from different geographical origins by comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry (GC × GC-TOF/MS) in combination with multivariate analysis[J]. J. Pharm. Biomed. Anal., 2007, 43: 1 721-1 727.