

(接 82 页后)

- [11] LEE K O, KIM S N, KIM Y C. Anti - wrinkle effects of water extracts of teas in hairless mouse [J]. *Toxicology Research*, 2014, 30: 283 - 289.
- [12] TREVITHICK J R, SHUM D T, REDAE S, et al. Reduction of sunburn damage to skin by topical application of vitamin E acetate following exposure to ultraviolet B radiation: effect of delaying application or of reducing concentration of vitamin E acetate applied [J]. *Scanning Microscopy*, 1993, 7: 1269 - 1281.
- [13] WU D C, GOLDMAN M P. A prospective, randomized, double - blind, split - face clinical trial comparing the efficacy of two topical human growth factors for the rejuvenation of the aging face [J]. *The Journal of Clinical and Aesthetic Dermatology*, 2017, 10: 31 - 35.
- [14] ELSNER P, FLUHR J W, GEHRING W, et al. Anti - aging data and support claims - consensus statement [J]. *Journal of the German Society of Dermatology*, 2011, 9: S1 - S32.
- [15] LEPHART E D. Skin aging and oxidative stress: Equol's anti - aging effects via biochemical and molecular mechanisms [J]. *Ageing Research Reviews*, 2016, 31: 36 - 54
- [16] SIKDAR S, PAPADOPOULOU M, DUBOIS J. What do we know about sulfuraphane protection against photoaging? [J]. *Journal of Cosmetic Dermatology*, 2016, 15: 72 - 77.
- [17] HSU S. Green tea and the skin [J]. *Journal of the American Academy of Dermatology*, 2005, 52: 1049 - 1059.
- [18] MEHTA R C, FITZPATRICK R E. Endogenous growth factors as cosmeceuticals [J]. *Dermatologic Therapy*, 2007, 20: 350 - 359.
- [19] ŻERA Ń SKA J, PASIKOWSKA M, SZCZEPANIK B, et al. A study of the activity and effectiveness of recombinant fibroblast growth factor (Q40P/S47I/H93G Rfgf - 1) in anti - aging treatment [J]. *Advances in Dermatology and Allergology*, 2016, 33: 28 - 36.
- [20] YOU J, ROH K B, LI Z, et al. The antiaging properties of *Andrographis paniculata* by activation epidermal cell stemness [J]. *Molecules*, 2015, 20: 17557 - 17569.
- [21] YOON B H, JEON Y H, HWANG B, et al. Anti - wrinkle effect of bone morphogenetic protein receptor 1a - extracellular domain (BMPR1a - ECD) [J]. *BMB Reports*, 2013, 46: 465 - 470.
- [22] KHAN T K, WENDER P A, ALKON D L. Bryostatins and its synthetic analog, picolog rescue dermal fibroblasts from prolonged stress and contribute to survival and rejuvenation of human skin equivalents [J]. *Journal of Cellular Physiology*, 2017. doi: 10.1002/jcp.26043.
- [23] TAGAMI H. Functional characteristics of the stratum corneum in photoaged skin in comparison with those found in intrinsic aging. *Archives of Dermatological Research*, 2008, 300: S1 - 6.
- [24] BISSETT D L, OBLONG J E, SAUD A. Topical niacinamide provides skin aging appearance benefits while enhancing barrier function [J]. *Journal of Clinical Dermatology*, 2003, 32S: 9 - 18.
- [25] LEVEQUE J L, CORCUFF P, De RIGAL J, et al. In vivo studies of the evolution of physical properties of the human skin with age [J]. *International Journal of Dermatology*, 1984, 23: 322 - 329.
- [26] 冯冰, 何聪芬, 赵华, 等. 生物技术在提高化妆品功效及功效检测中的应用 [J]. *日用化学品科学*, 2006, 29 (9) : 29 - 32.
- [27] ROBERTS W. Periorbital hyperpigmentation: review of etiology, medical evaluation and aesthetic treatment [J]. *Journal of Drugs in Dermatology*, 2014, 13: 472 - 482.
- [28] KLINGNAN D. Cosmeceuticals [J]. *Dermatologic Clinics*, 2000, 18: 609 - 615.
- [29] ANTONOPOULOU I, VARRIALE S, TOPAKAS E, et al. Enzymatic synthesis of bioactive compounds with high potential for cosmeceutical application [J]. *Applied Microbiology and Biotechnology*, 2016, 100: 6519 - 6543.
- [30] LEPHART E D, ACERSON M J, ANDRUS M B. Synthesis and skin gene analysis of 4' - acetoxy - resveratrol (4AR), therapeutic potential for dermal applications [J]. *Bioorganic & Medicinal Chemistry Letters*, 2016, 26: 3258 - 3262.
- [31] TAOFIQ O, GONZÁLEZ - PARAMÁS A M, BARREIRO M F, et al. Hydroxycinnamic acids and their derivatives: Cosmeceutical significance, challenges and future perspectives, a review [J]. *Molecules*, 2017, 22: 281.
- [32] SCHWARZ C, MEHNERT W, LUCKS J S, et al. Solid lipid nanoparticles (SLN) for controlled drug delivery. I. Production, characterization and sterilization [J]. *Journal of Controlled Release*, 1994, 30: 83 - 96.
- [33] NICHOLS J A, KATIYAR S K. Skin photoprotection by natural polyphenols: anti - inflammatory, antioxidant and DNA repair mechanisms [J]. *Archives of Dermatological Research*, 2010, 302: 71 - 83.
- [34] JIMÉNEZ Z, KIM Y J, MATHIYALAGAN R, et al. Assessment of radical scavenging, whitening and moisture retention activities of *Panax ginseng* berry mediated gold nanoparticles as safe and efficient novel cosmetic material [J]. *Artificial Cells, Nanomedicine, and Biotechnology*, 2017. doi: 10.1080/21691401.2017.1307216.
- [35] AZEEM A, RIZWAN M, AHMAD F J, et al. Emerging role of microemulsions in cosmetics [J]. *Recent Patents on Drug Delivery & Formulation*, 2008, 2: 275 - 289.
- [36] XAVIER - JUNIOR F H, VAUTHIER C, MORAIS A R, et al. Microemulsion systems containing bioactive natural oils: an overview on the state of the art [J]. *Drug Development and Industrial Pharmacy*, 2017, 43: 700 - 714.
- [37] United States. Food and Drug Administration. Guidance for Industry Considering Whether an FDA - Regulated Product Involves the Application of Nanotechnology [S]. 2014. <http://www.fda.gov/RegulatoryInformation/Guidances/ucm257698.htm>.
- [38] PEREIRA T A, GUERREIRO C M, MARUNO M, et al. Exotic vegetable oils for cosmetic O/W nanoemulsions: In vivo Evaluation [J]. *Molecules*, 2016, 21: 248.
- [39] Da ROCHA - FILHO P A, MARUNO M, FERRARI M, et al. Liquid crystal formation from sunflower oil: Long term stability studies [J]. *Molecules*, 2016, 21: 680.